

Recent Beach Progradation and High Resolution Ground Penetrating Radar Lines

Harry M. Jol, Department of Geography, University of Wisconsin - Eau Claire, 105
Garfield Ave., Eau Claire, WI; 54702, USA., jolhm@uwedc.edu

George Kaminsky, Washington Department of Ecology, P.O. Box 47690, Olympia, WA,
98504-7690, USA, gkam461@ecy.wa.gov

Peter Ruggerio, Washington Department of Ecology, P.O. Box 47690, Olympia, WA,
98504-7690, USA, prug461@ecy.wa.gov

During the summer of 1998, three high resolution ground penetrating radar (GPR) profiles were collected in coordination with global position system (GPS) data. A line was run in each of 3 subcells located within the larger Columbia River Littoral Cell – Ocean Shores (North Grays Harbor), Warrenton (South Grays Harbor) and Long Beach. These profiles were shot using 200 MHz antennae with 0.5 m step to allow for detailed imaging (<0.5 m resolution) of the subsurface. The processed profiles were topographically corrected using the GPS data and at each location, a common mid-point survey was collected to calculate velocity of the subsurface material so that depth calculations could be made. The data was collected in areas that the geographic information systems database reveals constant progradation over the time period of available topographic information. Further processing and interpretation, will aid in understanding rate of recent progradation and possible anomalies.




Figure 1 *Upper:* 100 MHz GPR profile (W-E) shot to show dipping foresets at Ocean Shores. *Lower:* 200 MHz GPR profile shot over same site showing higher resolution stratigraphy. The 1884 shoreline is located at east end of the profile.




Figure 2 200 MHz GPR profile (W-E) shot in 1998 along Warrenton-Cannery Road in the South Grays Harbor Cell. *Upper:* 1200 yr BP scarp (placer) is located at 20-40 m, *Middle:* 300 Yr BP scarp (placer) is located at 385-400 m, *Lower:* Further west along the line is an erosional scarp located at 585 – 606 m. The feature is in front of the last known earthquake event and could possibly be the 1886 El Nino event.




Figure 3 200 MHz GPR profile (W-E) shot in 1998 on the Long Beach Peninsula. *Upper:* 300 Yr BP scarp (placer) is located at 40 – 60 m. The 1873 shoreline is located at the same location. *Lower:* Further west along the line dunes are apparent in the topography as well as salt-water intrusion attenuates the radar signal. 1924 shoreline is located at approximately position 540.

Acknowledgements

The major support for this project comes from the Southwest Washington Coastal Erosion Study - United States Geological Survey (USGS). Further support comes from Sensors and Software, University of Wisconsin-Eau Claire, Portland State University, University College of the Fraser Valley and Grays Harbor College. We would also like to acknowledge the support of the Washington Department of Ecology (DOE), and Grays Harbor, Pacific, and Clatsop counties. Able bodied field assistance was provided by Brian Thayer. We thank Rilea Armed Forces Training Center, the Cranberry Research Foundation, and the communities within the study area for their aid in this project.

References

- Jol, H.M., Peterson, C.D., Vanderburgh, S. and Phipps, J. 1998. GPR as a regional geomorphic tool: shoreline accretion/erosion along the Columbia river littoral cell. Proceedings of the Seventh International Conference on Ground Penetrating Radar

(GPR'98), May 27-30, University of Kansas, Lawrence, KS, USA, Volume 1, pp. 257-262.

Jol, H.M., D.G. Smith, and R.A. Meyers, 1996. Digital ground penetrating radar (GPR): An improved and very effective geophysical tool for studying modern coastal barriers (examples for the Atlantic, Gulf and Pacific coasts, U.S.A.). *Journal of Coastal Research*, 12(4): 960-968.

Kaminsky, G.M., P. Ruggiero, G. Gelfenbaum, and C. Peterson, 1997. Long term coastal evolution and regional dynamics of a US Pacific Northwest Littoral Cell, *Coastal Dynamics*'97.

Meyers, R., D.G. Smith, H.M. Jol, and C.R. Peterson, 1996. Evidence for eight great earthquake-subsidence events detected with ground-penetrating radar, Willapa barrier, Washington. *Geological Society of America Geology*, 24: 99-102.

Phipps, J.B., 1990. Coastal accretion and erosion in Washington State 1977-1987 Shorelands and Coastal Zone Management Program. Washington Department of Ecology, Olympia, Washington.